

## Regular Article

# Changing parental depression and sensitivity: Randomized clinical trial of ABC's effectiveness in the community

Laura Perrone<sup>1</sup> , Steven D. Imrisek<sup>1,2</sup>, Allison Dash<sup>1,2</sup>, Melanie Rodriguez<sup>1,2</sup>, Erasma Monticciolo<sup>2</sup> and Kristin Bernard<sup>1</sup>

<sup>1</sup>Department of Psychology, Stony Brook University, Stony Brook, NY, USA and <sup>2</sup>Power of Two, Brooklyn, NY, USA

### Abstract

Attachment and Biobehavioral Catch-up (ABC) demonstrates efficacy in improving parent and child outcomes, with preliminary evidence for effectiveness in community settings. The objective of this study was to assess the effectiveness of a community-based ABC implementation in improving parent outcomes as well as to examine potential mediators and moderators of intervention effectiveness. Two hundred parents and their 5- to 21-month-old infants recruited from an urban community were randomly assigned to receive ABC or be placed on a waitlist. The majority of participants had a minority racial or ethnic background. Before intervention, parents completed questionnaires about sociodemographic risk and adverse childhood experiences. At both baseline and follow-up, parents reported depression symptoms and were video-recorded interacting with their infant, which was coded for sensitivity. The ABC intervention predicted significant increases in parental sensitivity and, among parents who completed the intervention, significant decreases in depression symptoms. Changes in parental depression symptoms did not significantly mediate the intervention effects on sensitivity. Risk variables did not moderate the intervention effects. The results indicate that ABC shows promise for improving parent outcomes in community settings, supporting dissemination.

**Keywords:** depression, early intervention, parenting, risk, sensitivity

(Received 17 September 2019; accepted 19 February 2020)

Preventive interventions in the first 5 years of life have been identified as essential both to improving the lives of children and families and to benefiting society more broadly (Doyle, Harmon, Heckman, & Tremblay, 2009). However, funding and providers for early interventions that enhance parenting are limited, making it important to ensure that resources are used for effective interventions. Attachment and Biobehavioral Catch-up (ABC), an early parenting intervention for young children exposed to adversity, has been examined extensively in efficacy studies, providing empirical support for its positive influence on parent and child outcomes. A critical next step is replicating these positive results beyond the lab. Thus, the present study evaluated the effectiveness of ABC when implemented by community providers in a community setting.

### Parenting Interventions and Programs

A variety of treatment programs and preventive interventions targeting children at risk for neglect or maltreatment in early

childhood have received empirical support. For example, Child-Parent Psychotherapy, a relationship-based treatment for mothers and children ages 0–5 who have experienced traumas, adversities, or harmful parenting practices, has been shown to improve parental mental health symptoms, child traumatic stress symptoms, child behavior problems, attachment security, and subsequent positive peer relations, among other positive outcomes (Guild, Toth, Handley, Rogosch, & Cicchetti, 2017; Lieberman, Ippen, & Van Horn, 2006; Lieberman, Van Horn, & Ippen, 2005). In addition, Parent-Child Interaction Therapy, which provides in vivo coaching to help parents learn positive interaction and behavioral management strategies, has been found to improve maternal sensitivity, maternal stress, child abuse potential, and child behavior among mother-child dyads with a history of or high risk for child maltreatment (Thomas & Zimmer-Gembeck, 2011). On a broader community level, the Positive Parenting Program, a multi-level preventive intervention targeting social and emotional competence in children, has been associated with reduced child maltreatment, child maltreatment injuries, and child out-of-home placements (Prinz, Sanders, Shapiro, Whitaker, & Lutzker, 2009). Finally, prenatal and postpartum home visits by a nurse through the Nurse Family Partnership program have been associated with a reduced number of substantiated reports of maltreatment (Eckenrode et al., 2017). These provide only a few examples of successful parenting programs and interventions (see Petersen, Joseph, & Feit (2014) for a

**Author for Correspondence:** Dr. Kristin Bernard, Department of Psychology, Stony Brook University, Stony Brook, NY 11794-2500; E-mail: [kristin.bernard@stonybrook.edu](mailto:kristin.bernard@stonybrook.edu).

**Cite this article:** Perrone L, Imrisek SD, Dash A, Rodriguez M, Monticciolo E, Bernard K (2020). Changing parental depression and sensitivity: Randomized clinical trial of ABC's effectiveness in the community. *Development and Psychopathology* 1–15. <https://doi.org/10.1017/S0954579420000310>

more thorough review). Although each of these programs has demonstrated success in reducing risk for maltreatment or neglect, many families at risk currently do not receive adequate services (Petersen *et al.*, 2014). As a result, continued evaluation of prevention programs is necessary with the goal of identifying programs that can be sustainably and effectively implemented in the community.

### *ABC Intervention*

One program that shows promise for decreasing the gap between the need and availability of community preventive services is ABC. The ABC intervention is a 10-session home-visiting program designed to support infants in contexts of neglectful or problematic parenting (Dozier & Bernard, 2017, 2019). The intervention was initially developed for caregivers of infants in foster care and has since been expanded for use with parents with child protective services involvement, parents of toddlers with child welfare involvement, and parents of internationally adopted children (Dozier, Roben, Caron, Hoye, & Bernard, 2018). The intervention targets three specific parent behaviors to promote secure attachment and healthy biological regulation in children: (a) nurturance, or responsiveness to the child's distress; (b) following the child's lead, or responsiveness to the child when not distressed; and (c) reducing frightening behaviors.

The ABC intervention promotes these parenting behaviors through information provided to parents addressing each target and "in-the-moment" comments delivered by parent coaches during home visits. In-the-moment comments seek to describe the parent's behavior, link the behavior to an intervention target, and indicate the long-term effects of that behavior. Parent coaches' in-the-moment comments have been identified as an active ingredient promoting parental behavior change in ABC (Caron, Bernard, & Dozier, 2018). Consistent with meta-analytic findings (Bakermans-Kranenburg, van IJzendoorn, & Juffer, 2003), other important features of ABC that may contribute to its effectiveness include its short duration, targeted focus on changing parenting behavior, and implementation in the typical home environment (Dozier & Bernard, 2017).

### *ABC Intervention Efficacy*

Substantial evidence from randomized clinical trials supports the efficacy of ABC in increasing positive parenting behaviors. For example, foster mothers who received ABC intervention showed greater increases from preintervention to postintervention in observed sensitivity toward their infant than foster mothers who received a control intervention matched for duration and structure (Bick & Dozier, 2013). Similarly, mothers from low income communities who received ABC showed greater increases in sensitivity and greater decreases in intrusiveness from preintervention to postintervention than mothers who received a control intervention (Yarger, Hoye, & Dozier, 2016). In addition, parents of internationally adopted children who received ABC demonstrated greater postintervention sensitivity and positive regard than adoptive parents who received a control intervention, with differences remaining 1.5–2.5 years postintervention (Yarger, Bernard, Caron, Wallin, & Dozier, 2019). Therefore, a strong evidence base supports ABC as an intervention for increasing parental sensitivity.

The benefits of ABC extend beyond parent behavior changes to a wide range of child outcomes. Among foster children, ABC as

compared with a control intervention has been associated with higher rates of attachment security (Dozier *et al.*, 2009), more typical diurnal cortisol patterns (Dozier *et al.*, 2006), higher receptive vocabulary (Bernard, Lee, & Dozier, 2017), and greater executive functioning (Lewis-Morrarty, Dozier, Bernard, Terracciano, & Moore, 2012). Similarly, foster children who received the ABC version for toddlers (ABC-T) as compared with a control intervention displayed higher postintervention receptive vocabulary (Raby, Freedman, Yarger, Lind, & Dozier, 2019) and executive functioning (Lind, Raby, Caron, Roben, & Dozier, 2017). Among children of parents who are involved with child protective services, receiving ABC was associated with higher rates of attachment security (Bernard *et al.*, 2012), more typical diurnal cortisol patterns (Bernard, Dozier, Bick, & Gordon, 2015; Bernard, Hostinar, & Dozier, 2015), greater child compliance (Lind, Bernard, Yarger, & Dozier, 2019), and less child negative affect during a challenge (Lind, Bernard, Ross, & Dozier, 2014) than receiving a control intervention was. Follow-up studies have also shown effects on respiratory sinus arrhythmia, heart rate, and more normative patterns of brain activity in middle childhood (Bick, Palmwood, Zajac, Simons, & Dozier, 2019; Tabachnick, Raby, Goldstein, Zajac, & Dozier, 2019). Taken together, these studies support the efficacy of ABC in improving both parent and child outcomes.

### *ABC Intervention Effectiveness*

Given the strong evidence of ABC's efficacy, a crucial next step is examining ABC's effectiveness when implemented in community-based settings. Intervention efficacy within a research context does not always guarantee parallel effectiveness in applied practice (Glasgow, Lichtenstein, & Marcus, 2003; Weisz, Weiss, & Donenberg, 1992). The National Institute of Mental Health has therefore called for increased efforts to conduct effectiveness studies testing interventions beyond the laboratory (Hoagwood & Olin, 2002). Important current issues within effectiveness research include dissemination of interventions to diverse populations and identification of mediators and moderators of treatment success (La Greca, Silverman, & Lochman, 2009), each of which are relevant to the further development of ABC.

To date, a few studies have examined the effectiveness of implementing ABC in community-based settings using pre-post designs. In one of the earliest examples, ABC was disseminated in a diverse Hawaiian community through a variety of organizations, with steps taken to ensure parent coach commitment and fidelity to the model while also allowing for relevant cultural modification (Caron, Weston-Lee, Haggerty, & Dozier, 2016). Caregivers who completed ABC in this sample demonstrated significant pre- to postintervention increases in observed sensitivity (*i.e.*, following the child's lead) and delight and decreases in observed intrusiveness.

The effectiveness of ABC is further supported by an examination of ABC dissemination across five community sites from July 2013 to May 2016 when implemented by 23 coaches with varying levels of educational training and diverse professional backgrounds (Roben, Dozier, Caron, & Bernard, 2017). Replicating the previous study, families who completed ABC demonstrated a significant increase in observed sensitivity from pre- to postintervention with a large effect size. It is important to note that both of these studies reflect intraindividual changes, rather than interindividual differences between treatment and control groups. Nonetheless, the findings support ABC's effectiveness in diverse community settings.

In addition, ABC has been disseminated outside its lab of origin. For example, Berlin, Shanahan, and Appleyard Carmody (2014) implemented ABC within a residential substance-abuse treatment program for mothers. Compared with mothers receiving a control “book-of-the-week” program, mothers receiving ABC demonstrated increased postintervention sensitive parenting behavior with a medium effect size. In addition, Berlin, Martoccio, and Jones Harden (2018) tested community implementation of ABC in a randomized controlled trial comparing Early Head Start (EHS) programs including ABC or a control book-of-the-week program. The results indicated that ABC was associated with greater postintervention sensitivity, lower intrusiveness, and greater positive regard. Furthermore, infants whose mothers received ABC displayed more rapid postintervention cortisol recovery to a mild stressor than those who received the book-of-the-week program (Berlin, Martoccio, Bryce, & Harden, 2019). These results offer evidence for ABC’s effectiveness in a predominantly Latino population and among low-income families that were not recruited for additional risk indicators, highlighting a preventive context. Taken together, these studies provide preliminary evidence for the effectiveness of ABC as it is disseminated in community settings. At the same time, additional effectiveness trials are necessary to replicate the current successes in more diverse community settings, particularly by using randomized clinical trials (RCTs).

Additional work is necessary to understand outcomes, mechanisms, and moderators of ABC. Although extensive research has supported ABC’s effects on parental behavior targets and child outcomes, fewer studies have investigated additional parent outcomes. Studies also have yet to examine possible mechanisms of ABC’s effectiveness. In-the-moment commenting has been identified as one active ingredient of ABC that promotes changes in parental sensitivity and intrusiveness (Caron et al., 2018); however, it is likely that there are other mechanisms of change occurring within the parent and parent–child dyad. Finally, few moderators that elucidate for whom ABC is most effective have been identified. When ABC was implemented along with Head Start in a population of low-income mothers, greater preintervention intrusiveness and maternal secure or anxious (rather than avoidant) attachment style predicted improved parent outcomes, highlighting the possibility that parent characteristics may moderate ABC outcomes (Berlin et al., 2018). Understanding additional parent outcomes, mediators, and moderators will help maximize the positive effects of ABC as the intervention moves from efficacy studies to effectiveness trials and dissemination efforts.

### *Identifying Mechanisms of Change in Sensitivity: The Role of Parental Depression*

Sensitivity, or responding contingently to children’s cues (Ainsworth, Bell, & Stayton, 1974), is an important predictor of developmental outcomes, including child attachment (De Wolff & van IJzendoorn, 1997; Zeegers, Colonesi, Stams, & Meins, 2017), physiological reactivity (Albers, Riksen-Walraven, Sweep, & de Weerth, 2008), socioemotional development (Leerkes, Blankson, & O’Brien, 2009), executive functioning (Valcan, Davis, & Pino-Pasternak, 2018), language development (Barnett, Gustafsson, Deng, Mills-Koonce, & Cox, 2012), and symptoms of psychopathology (Haltigan, Roisman, & Fraley, 2013). Moreover, the association between sensitivity and positive child outcomes holds in diverse populations, including among ethnic minority families (Mesman, van IJzendoorn, & Bakermans-Kranenburg, 2012). As described

above, increased parental sensitivity is a consistent outcome of ABC intervention, but less is known about whether changes in other parent outcomes may mediate these effects.

Current literature consistently supports an association between parental depression and sensitivity. In a meta-analysis of 46 observational studies from 1974 to 1996, maternal depression had a moderate association with increased disengaged and negative/hostile behavior toward children as well as a small association with fewer positive social interaction behaviors (Lovejoy, Graczyk, O’Hare, & Neuman, 2000). Similarly, in a review of studies over the following 10 years, Field (2010) found that depressed mothers show decreased sensitivity across cultures and socioeconomic statuses. More recently in a meta-analysis examining the first year of life, maternal depression was significantly associated with a small effect on observed maternal sensitivity (Bernard, Nissim, Vaccaro, Harris, & Lindhiem, 2018). Furthermore, paternal depression has also been associated with withdrawn paternal behavior in father–infant interactions at 3 months (Sethna, Murray, Netsi, Psychogiou, & Ramchandani, 2015).

The association between depression and sensitivity affords the possibility that changes in these variables may be related. Many studies have provided preliminary evidence that interventions with depressed caregivers do indeed promote changes in parental sensitivity.

A meta-analysis examining the effect of psychotherapy for depressed mothers indicated intervention results in decreased levels of maternal depression, improved mother–child interactions, and improved child mental health (Cuijpers, Weitz, Karyotaki, Garber, & Andersson, 2015). Similarly, in a review of RCTs assessing the effects of postnatal depression treatments, interventions were found generally to improve the quality of interaction between parent and child (Poobalan et al., 2007). Finally, a meta-analysis of preventive interventions for depressed mothers found a small-to-medium effect size on sensitivity, with interventions including baby massage or support groups being more effective than those with individual therapy (Kersten-Alvarez, Hosman, Riksen-Walraven, van Doesum, & Hoefnagels, 2011). It is noteworthy that interventions with a behavioral or support-based component had larger effects on parental sensitivity, both of which may be provided by ABC.

Given that interventions targeting mothers with postnatal depression frequently produce changes in sensitivity, changes in maternal depression may be one mechanism through which interventions promote improved sensitivity. Additionally, improving caregiver well-being has been identified as central to promoting positive outcomes in both caregivers and children, with the caregiver’s ability to self-regulate highlighted as essential to responding effectively to challenging situations (Luthar & Eisenberg, 2017). If ABC does produce improvements in caregivers’ depressive symptoms, changes in caregiver depression may serve as a mediator through which ABC improves caregiver sensitivity.

### *Examining Moderators of ABC Effectiveness*

#### *Cumulative risk*

In considering moderators that predict for whom ABC intervention is most successful, one potential variable is cumulative risk, or the additive effects of exposure to multiple risk factors. Cumulative risk has consistently been associated with less sensitive parenting (e.g., Lengua, Honorado, & Bush, 2007; Trentacosta et al., 2008), a primary outcome of ABC. However, the effects of cumulative risk on interventions targeting parenting is not known as current

literature provides mixed results. Among families of young children with developmental delay receiving Parent–Child Interaction Therapy, cumulative risk predicted dropout and diminished treatment effects (Bagner & Graziano, 2013), suggesting that risk may interfere with intervention success. In contrast, cumulative risk was associated with greater treatment effectiveness in families receiving high-intensity Parent Management Training (Tømmerås & Kjøbli, 2017), offering the possibility that families with higher risk have greater potential for change. Finally, cumulative risk did not predict differences in parenting outcomes for an intervention that was directed at positive parenting of toddlers and a family-focused preventive intervention for parents of adolescents (Spoth et al., 1998; Stolk et al., 2008). Given these varying findings, additional studies are necessary to elucidate the role of cumulative risk as a moderator of parenting intervention outcomes.

#### *Caregiver adverse childhood experiences*

Another risk factor that may moderate intervention effectiveness is the caregiver's level of adverse childhood experiences (ACEs). Childhood maltreatment, one category of ACEs, has been associated with less sensitivity, greater hostility, and greater child abuse potential in mothers (Bailey, DeOliveira, Wolfe, Evans, & Hartwick, 2012; Bert, Guner, Lanzi, & Centers for Prevention of Child Neglect, 2009; Pereira et al., 2012). Similarly, greater ACEs have been associated with greater likelihood of lifetime depressive disorders (Chapman et al., 2004), which in turn has been associated with parental sensitivity as described above. However, the influence of ACEs on intervention effectiveness has not been examined. Parents with ACEs may stand to benefit the most from intervention given the association between ACEs and decreased sensitivity. At the same time, caregivers' ACEs may interfere with their ability to benefit from intervention. Examining cumulative risk and parental ACEs as potential moderators of ABC's influence on parent outcomes will provide insight into which parents are most likely to benefit from ABC.

#### *The Present Study*

Using an RCT in a community setting, the present study attempted to replicate previous findings that ABC increases sensitive parental behavior. In addition, we examined changes in parental depression as a potential mechanism for increased sensitivity and measures of risk as potential moderators of intervention effectiveness. We hypothesized that parents who received ABC through the community implementation would show greater improvements in parental sensitivity than parents in a waitlist control condition. In addition, we expected that ABC would reduce parents' depressive symptoms, which would serve as a mechanism through which ABC improves parental sensitivity. Finally, given that cumulative risk and parental ACEs have the potential to interfere with intervention effectiveness or to provide greater opportunities for intervention-promoted change, we conducted exploratory analyses to examine these variables as potential moderators of intervention effectiveness on parent outcomes.

#### **Method**

##### *Implementation of ABC*

The current study was conducted in collaboration with Power of Two, a nonprofit organization in Brooklyn, New York. Power of Two was founded in 2015 to scale the dissemination of ABC to

families living in poverty and involved in the child welfare system in New York City. The RCT reported here results from an evaluation of the effectiveness of implementing ABC through Power of Two. Across the study period, ABC was provided by 24 parent coaches who varied in their racial/ethnic, professional, and educational backgrounds. Six (25%) of the coaches were Caribbean, 6 (25%) were Hispanic or Latin, 5 (20.83%) were African American, 3 (12.5%) were Caucasian, 2 (8.3%) were biracial/multiethnic, and 2 (8.3%) identified as other. In terms of education, 4 (16.7%) of the coaches had a high school degree, 1 (4.2%) had completed some college, 12 (50%) had a bachelor's degree, and 7 (29.2%) had a master's degree. Twenty-three (95.8%) had previous experience working with families or children, 12 (50%) had previous social work experience, and 7 (29.2%) had previous clinical work experience.

Suitability to be a parent coach was determined regardless of professional and educational background and based on selection criteria defined by the model developer: (a) evidence of valuing attachment/openness (as assessed via an abbreviated attachment interview) and (b) initial skill in delivering in-the-moment feedback on parent–child interactions (as assessed via a video vignette-based role-play; Caron, Roben, Yarger, & Dozier, 2018). Parent coaches participated in a 2-day training followed by a year of weekly supervision provided by the model developer's team. Weekly supervision included 1 hr of clinical group supervision focused on case conceptualization and 30 min of individual fidelity-focused supervision (for a description of the standardized training and supervision practices, see Caron et al., 2016). Fidelity was monitored by having the parent coach and a reliable ABC fidelity coder randomly code selected 5-min segments of the coach's sessions for on-target parental behaviors and the coach's rate and accuracy of commenting on such behaviors. The parent coach and fidelity coder met weekly to discuss the accuracy of coding as well as the parent coach's frequency and content of in-the-moment comments (for more details on fidelity monitoring and related outcomes, see Caron & Dozier, 2019).

#### *Participants*

The participants were 200 primary caregivers (referred to as parents throughout text) and children from a large urban area on the east coast. A power analysis conducted using G\*Power 3.1.7 demonstrated that a sample size of 134 would afford 90% power to detect an effect size between small and medium ( $f^2 = .08$ ) by using linear multiple regression with baseline sensitivity as a covariate. Estimating approximately 30% attrition based on similar studies, we planned to recruit 200 participants in total. At the initial intake visit, children ranged in age from 5.64 to 21.48 months ( $M = 11.82$ ,  $SD = 4.36$ ). Ninety-eight (49.0%) of the children were male. One hundred twenty-three (61.5%) of the children were African American, 32 (16.0%) were biracial/multi-ethnic, 28 (14.0%) were Hispanic or Latin, 8 (4.0%) were Caribbean, 2 (1.0%) were American Indian or Alaska Native, 1 (0.5%) was Caucasian, 5 (2.5%) identified as other, and 1 (0.5%) did not report race/ethnicity information.

At the initial intake visit, parents ranged in age from 17.04 to 64.89 years ( $M = 29.85$ ,  $SD = 7.32$ ). Eight (4.0%) of the parents were male. One hundred eighty-three (91.5%) of the parents in this study were the child's birth parent. Of parents who provided their household composition, 65.8% lived without a partner and 40.8% lived alone or only with their children/grandchildren. The majority of the sample experienced high housing risk, with

35.9% of parents providing housing information reporting living in a shelter and 25.5% reporting living in public housing developments. One hundred twenty-nine (64.5%) of the parents were African American, 30 (15.0%) were Hispanic or Latin, 17 (8.5%) were biracial/multi-ethnic, 10 (5.0%) were Caribbean, 2 (1.0%) were American Indian or Alaska Native, 1 (0.5%) was Asian, 3 (1.5%) were Caucasian, 5 (2.5%) identified as other, and 3 (1.5%) did not report race/ethnicity information. The majority (83.0%) of parents who responded to a question about monthly income reported an income level below the poverty threshold for the number of children and adults in their household.

### Procedure

Power of Two outreach staff recruited parents and children from the community to participate in the study through partnerships with community-based organizations (e.g., family-based homeless shelters, community fairs, health service providers) and by talking directly with parents in the neighborhood (e.g., on the street, in public housing developments). Participants were recruited between January 2016 and November 2018 and were drawn from the population served by Power of Two. Upon contacting parents interested in receiving ABC, Power of Two staff asked about interest in participating in a research study. If the parent expressed interest, they were contacted by the research coordinator, who provided additional information about the study and screened for eligibility. Inclusion criteria for the parent included being at least 16 years old and having custody of a child between 6 and 20 months of age. Exclusion criteria included significant child medical or developmental delay, not living in the service area, and not speaking English. See the CONSORT diagram in [Figure 1](#) for the number of potential dyads considered at each stage.

The research coordinator attended the initial intake appointment of eligible dyads to describe the study in detail, answer questions, and ensure parental understanding. If the parent was 18 years of age or older, the research coordinator obtained informed consent from the parent. If the parent was under 18 years of age, consent was obtained from both the parent and the parent's legal guardian. All dyads then completed the initial intake typically provided as part of Power of Two's services, which included completion of questionnaires and a video-recorded parent-child interaction.

After providing informed consent, dyads were randomly assigned to the ABC or the waitlist control condition by the principal investigator using a randomly generated number sequence, allowing the research staff to remain masked to condition prior to randomization. Given that research staff within Power of Two were responsible for coordinating between parents and parent coaches, they were aware of random assignment thereafter. Those assigned to ABC began the intervention immediately, with home visits scheduled at approximately weekly intervals. Among dyads that completed all 10 sessions of ABC, the average number of weeks to completion was 14.39 ( $SD = 3.84$ ). Dyads assigned to the waitlist condition did not receive any intervention during a waiting period. After the intervention or waiting period, dyads completed a follow-up visit at the Power of Two office in Central Brooklyn between June 2016 and September 2018. The average time between baseline and follow-up visits was 7.19 months ( $SD = 2.85$ ). An attempt was made to conduct a follow-up assessment for all dyads. During the follow-up visit, the parent

completed questionnaires and repeated the video-recorded parent-child interaction. The parent received \$50 for participating in the follow-up research visit. Dyads assigned to the waitlist control were then provided the opportunity to complete ABC.

### ABC Intervention

The ABC intervention consists of 10 weekly home visits (for a detailed description of the intervention, see Dozier & Bernard, 2017). In these sessions, the parent coach discusses the rationale and evidence supporting each ABC target: nurturance (sessions 1 and 2), following the lead (3 and 4), and reducing frightening behaviors (5 and 6). In later sessions, the parent coach helps the parent consider how their own upbringing and/or current situation may interfere with providing sensitive care (sessions 7 and 8). In the final two sessions (9 and 10), the parent coach reviews previous content and celebrates progress. Across all sessions, the parent coach makes frequent in-the-moment comments on parent-child interactions promoting the behavior targets during both spontaneous interactions and structured activities with the child. Video feedback is also used to further reinforce the parent's progress and highlight successes.


### Measures

#### Parental behavior

Parental behavior was assessed through a semistructured, video-recorded play interaction in which the child was placed in an infant seat and provided with developmentally appropriate toys. The parent was instructed to interact with their child as they normally would, first from a distance without touching the child or toys for 2 min and then as close as they would like for 7 min. Parental behavior was coded on 5-point scales adapted from the Qualitative Scales of the Observational Record of the Caregiving Environment (National Institute of Child Health and Human Development Early Child Care Research Network, 1999, 2003) for "sensitivity" (following the child's lead), which reflected the parent's ability to respond contingently to the child's cues and behaviors; "intrusiveness," which reflected the parent's tendency to direct or overstimulate the child; and "positive regard," which reflected the parent's expression of enjoyment of the child. These scales have been used previously to assess parental behavior in samples with a similar age range and demographic diversity (e.g., Lind et al., 2019). Each play interaction was double-coded by trained coders who were masked to the dyad's assigned condition. Raters showed good reliability, with baseline and follow-up one-way average measures intraclass correlation coefficients of .62 and .79 for sensitivity, .79 and .84 for intrusiveness, and .75 and .74 for positive regard. See [Table 1](#) for means and standard deviations.

#### Parental depression

Parental depression was assessed with the Center for Epidemiological Studies Depression Scale (CES-D; Radloff, 1977). This 20-item self-report scale assesses frequency of depression symptoms (e.g., "I felt lonely" and "I had crying spells") over the previous week on a 4-point scale ranging from 0 = "rarely or none of the time (less than 1 day)" to 3 = "most or all of the time (5-7 days)." The CES-D has been found to have strong psychometric properties across diverse adult populations with reliabilities ranging from .85 to .87 in different racial/ethnic groups (Radloff, 1977; Roberts, 1980). Scores ranged from 0 to



**Figure 1.** Participant flow diagram for this randomized clinical trial of the ABC intervention.

**Table 1.** Means, standard deviations, and bivariate correlations of primary variables

Variable	Mean (SD)	1	2	3	4	5	6	7	8	9
1. Baseline sensitivity	2.38 (0.73)	–								
2. Baseline intrusiveness	3.18 (0.99)	–.55***	–							
3. Baseline positive regard	3.46 (0.77)	.54***	–.05	–						
4. Baseline depression	14.56 (10.56)	–.00	.00	.01	–					
5. Follow-up sensitivity	2.84 (0.83)	.35***	–.30***	.20*	–.22**	–				
6. Follow-up intrusiveness	2.37 (1.00)	–.23**	.38***	.12	.15	–.50***	–			
7. Follow-up positive regard	3.47 (0.77)	.27**	–.16	.35***	–.12	.62***	–.24**	–		
8. Follow-up depression	14.73 (11.30)	–.22**	.04	–.20*	.57***	–.36***	.15	–.27**	–	
9. Baseline ACE score	3.97 (2.75)	–.15*	.09	–.04	.40***	–.18*	.15	–.09	.44***	–
10. Cumulative risk	3.12 (1.16)	–.06	.01	–.02	.12	–.18*	.24**	–.13	.12	.12

Note: \* $p < .05$  \*\* $p < .01$ . \*\*\* $p < .001$ .

53 at baseline and from 0 to 46 at follow-up, with 36.9% and 38.8% of parents scoring at or above the clinical cutoff of 16 at baseline and follow-up, respectively. The Cronbach coefficient alphas for baseline and follow-up CES-D in the present study were .75 and .79, respectively. See Table 1 for means and standard deviations.

#### Adverse childhood experiences

ACEs were assessed with the ACE questionnaire, which asks whether the parent experienced each of the following adverse events during childhood: emotional abuse, physical abuse, sexual abuse, emotional neglect, physical neglect, single-parent home, witnessing domestic violence, substance-abusing parent, incarcerated parent, or parent with psychopathology (Felitti et al., 1998). Retrospective reports of ACE scores by adults have been shown to have good test-retest reliability with kappa of .64 and low rates of discordance (Dube, Williamson, Thompson, Felitti, & Anda, 2004). Scores at intake ranged from 0 to 10 in the present sample ( $M = 3.97$ ,  $SD = 2.75$ ), with 51.3% of parents reporting a score of 4 or more.

#### Cumulative risk

On questionnaires, parents were asked to report on demographic factors used to assess risk. A cumulative risk score was formed based on the following risk factors: single parent (63.2%), housing risk (living in a shelter; 33.8%), young parent (age 21 or younger at child's birth; 16.0%), parental involvement with the child welfare system (40.4%), and low parental education (less than high school or equivalent degree; 31.6%). Two risk factors were not included in our cumulative risk score because they were present in almost all dyads: financial risk (receiving some form of assistance such as food stamps or having a household income that falls below the poverty threshold; 99.4%) and child racial/ethnic minority status (99.5%). Dyads received one point for each risk indicator present. Missing scores on individual risk items were replaced with the mean for that risk item. Cumulative risk scores ranged from 0.99 to 6.0 ( $M = 3.12$ ,  $SD = 1.16$ ).

#### Analyses

As a preliminary analysis, we conducted a randomization check using  $t$  tests and chi-square tests to determine whether there were any baseline differences in demographic, risk, or primary

outcome variables between participants assigned to the ABC and waitlist control conditions. We conducted the same comparison analyses within the ABC group to determine whether there were any differences between those who did and did not complete all 10 sessions of the intervention.

For our principal analyses, we analyzed the data by using structural equation modeling with robust maximum likelihood estimation in Mplus 8 Version 1.5 (Muthén & Muthén, 2012–2017). We conducted cross-lagged analyses to examine whether participation in ABC as opposed to the waitlist control predicted follow-up parental behavior and depression scores accounting for baseline scores and child and parent ages. Initial more conservative analyses included all participants and used all available scores. Additional analyses were conducted excluding dyads assigned to the ABC group who had dropped out of the intervention before completing all 10 sessions. In addition, we conducted a mediation analysis within this cross-lagged model to examine whether follow-up depression served as a mechanism through which ABC influenced follow-up sensitivity. Finally, we conducted a series of exploratory moderation analyses in PROCESS 2.13 (Hayes, 2014) to examine whether risk variables (ACE and cumulative risk scores) served as moderators of the association between the ABC intervention and significant follow-up variables.


## Results

### Preliminary Analyses

Means, standard deviations, and bivariate correlations between primary variables are found in Table 1.

### Randomization check

There were no significant differences between the ABC and waitlist control groups on demographic variables including child gender, child or parent racial/ethnic minority status, parent or child age at intake, and whether the parent was the child's birth parent. In addition, the groups did not significantly differ on risk variables including baseline ACE and cumulative risk. Finally, the groups did not significantly differ on baseline levels of the primary variables examined in analyses including parental sensitivity, intrusiveness, positive regard, and depression symptoms.



**Figure 2.** Path diagram with standardized coefficients for structural equation cross-lagged model of parental sensitivity and parental depression symptoms for full sample. Model fit statistics indicated good fit: RMSEA = .00, CFI = 1.00, and TLI = 1.02. Nonsignificant covariances between age covariates and baseline and follow-up variables not depicted for simplicity. † $p < .10$ . \* $p < .05$ . \*\* $p < .01$ . \*\*\* $p < .001$ .

### Dropout analyses

Of the 100 dyads randomly assigned to receive ABC, 37 (37%) did not complete all 10 sessions of the intervention, with 15 completing no sessions, 7 completing 1 session, 4 completing 2 sessions, 1 completing 3 sessions, 5 completing 4 sessions, 4 completing 7 sessions, and 1 completing 8 sessions (see Supplementary Table 1 for descriptive information). Dyads who did or did not complete all 10 sessions of ABC did not differ on the demographic variables of child gender, child or parent racial/ethnic minority status, parent or child age at intake, and whether the parent was the child's birth parent. Parents who did not complete all 10 sessions had lower baseline scores for sensitivity than parents who completed all 10 sessions,  $t(89) = -2.02$ ,  $p = .047$ , but did not differ in baseline intrusiveness, positive regard, or depression scores. Parents who did not complete all 10 sessions also had significantly higher cumulative risk scores,  $t(98) = 2.86$ ,  $p = .005$ , but did not differ in ACE score.

### Changes in Parental Behavior and Depression Symptoms

#### Full sample analysis

First, we examined whether dyads randomly assigned to participation in ABC displayed greater follow-up observed parental behavior and lower depression symptoms than the waitlist control group by using a cross-lagged structural equation model that accounted for baseline parental behavior, baseline parental depression symptoms, and child and caretaker age at intake. A separate model was run for each parental behavior. Figure 2 shows the cross-lagged model examining ABC's influence on parental sensitivity and depression symptoms, with unstandardized coefficients and significance levels. The model fit statistics, root mean square error of approximation (RMSEA) = 0.00, comparative fit index (CFI) = 1.00, and Tucker-Lewis index (TLI) = 1.02, were within the range recommended for good model fit:  $RMSEA \leq 0.06$  and  $CFI/TLI \geq 0.95$  (Hu & Bentler, 1999). The model indicated that assignment to the ABC group was associated higher follow-up sensitivity,  $b = 0.26$ ,  $p = .035$  (see Table 2 and Figure 2). Although not significant, assignment to the ABC group was associated with lower follow-up depression symptoms,  $b = -2.77$ ,  $p$

$= .062$ . Using a last observation carried forward approach, on average parents who were assigned to the intervention condition had a greater increase in sensitivity ( $M = 0.41$ ,  $SD = 0.83$ ) than parents who were assigned to the waitlist group ( $M = 0.24$ ,  $SD = 0.80$ ). The effect size for these mean differences was small, Cohen  $d = 0.21$ , 95% CI [-0.08, 0.50]. In contrast, ABC did not have a significant effect on follow-up levels of intrusiveness and positive regard in the cross-lagged model corresponding to each.

#### ABC intervention completers


When the cross-lagged models were repeated comparing only dyads who completed all 10 sessions of ABC and the waitlist group, a similar pattern of results emerged. Figure 3 shows the cross-lagged model examining ABC's influence on parental sensitivity and depression symptoms with unstandardized coefficients and significance levels. The model fit statistics indicated good model fit, RMSEA = 0.05, CFI = 0.98, and TLI = 0.96. The model indicated that assignment to the ABC group was associated with higher follow-up sensitivity,  $b = 0.35$ ,  $p = .007$ , and lower follow-up depression symptoms,  $b = -3.76$ ,  $p = .013$  (see Table 3 and Figure 3). Using a last observation carried forward approach, on average parents who completed ABC displayed greater increases in sensitivity ( $M = 0.60$ ,  $SD = 0.88$ ) and greater decreases in depression symptoms ( $M = -1.65$ ,  $SD = 7.65$ ) than parents assigned to the waitlist condition ( $M = 0.24$ ,  $SD = 0.80$  for sensitivity;  $M = 0.46$ ,  $SD = 9.28$  for depression symptoms). The effect sizes corresponding to these mean differences were small-to-moderate for parental sensitivity, Cohen  $d = 0.43$ , 95% CI [0.10, 0.76], and small for parental depression symptoms, Cohen  $d = 0.24$ , 95% CI [0.08, 0.56]. However, completion of ABC did not significantly affect follow-up levels of intrusiveness and positive regard.

#### Parental Depression Mediating ABC Effects on Parental Sensitivity

Given the direct effects reported above, we next examined whether follow-up parental depression symptoms mediated ABC effects on parenting behavior, focusing specifically on parental sensitivity for ABC completers versus waitlist participants.

**Table 2.** Model estimated parameters for the full-sample structural cross-lagged model

Effect	Unstandardized Estimate	SE	Est/SE	p	95% CI	
					Lower	Upper
Follow-up sensitivity ON						
Baseline sensitivity	0.38	0.09	4.34	.00	0.21	0.55
Baseline depression symptoms	-0.02	0.01	-2.73	.01	-0.03	-0.00
Caregiver age at intake	0.01	0.01	0.61	.54	-0.01	0.02
Child age at intake	-0.14	0.18	-0.79	.43	-0.49	0.21
Intervention group	0.26	0.12	2.11	.04	0.02	0.50
Follow-up depression ON						
Baseline sensitivity	-3.18	1.08	-2.94	.00	-5.30	-1.06
Baseline depression symptoms	0.59	0.07	8.47	.00	0.45	0.73
Caregiver age at intake	0.10	0.10	1.01	.31	-0.10	0.30
Child age at intake	-0.48	2.13	-0.22	.82	-4.66	3.70
Intervention group	-2.77	1.48	-1.87	.06	-5.68	0.13
Baseline sensitivity WITH						
Baseline depression symptoms	0.13	0.59	0.22	.83	-1.02	1.28
Follow-up sensitivity WITH						
Follow-up depression symptoms	-1.43	0.55	-2.57	.01	-2.51	-0.34
Caregiver age at intake WITH						
Baseline sensitivity	1.25	0.43	2.90	.00	0.41	2.10
Baseline depression symptoms	-1.22	5.84	-0.21	.83	-12.66	10.22

**Figure 3.** Path diagram with standardized coefficients for structural equation cross-lagged model of parental sensitivity and parental depression symptoms for completers. Nonsignificant covariances between age covariates and baseline and follow-up variables not depicted for simplicity. Model fit statistics indicated good fit: RMSEA = .05, CFI = .98, and TLI = .96. \* $p < .05$ . \*\* $p < .01$ . \*\*\* $p < .001$ .

Mediation analyses were conducted in MPlus within the cross-lagged model reported above. The model fit statistics indicated good fit, RMSEA = 0.05, CFI = 0.98, and TLI = 0.96. The total indirect effect, although in the expected direction, was not significant,  $b = 0.07$ ,  $p = .084$  (see Supplementary Table 2).

As an additional follow-up analysis to examine the directionality of any potential mediation effect, we repeated the same analysis with parental depression symptoms as the outcome and parental sensitivity as the mediator. The model fit statistics indicated good fit, RMSEA = 0.05, CFI = 0.98, and TLI = 0.96. The

**Table 3.** Model estimated parameters for the completer-only structural cross-lagged model

Effect	Unstandardized estimate	SE	Est/SE	p	95% CI	
					Lower	Upper
Follow-up sensitivity ON						
Baseline sensitivity	0.40	0.09	4.38	.00	0.22	0.58
Baseline depression symptoms	-0.02	0.01	-2.71	.01	-0.03	-0.00
Caregiver age at intake	0.01	0.01	0.48	.63	-0.01	0.02
Child age at intake	-0.11	0.19	-0.58	.56	-0.48	0.26
Intervention group	0.35	0.13	2.69	.01	0.10	0.60
Follow-up depression ON						
Baseline sensitivity	-2.48	1.10	-2.25	.02	-4.64	-0.32
Baseline depression symptoms	0.59	0.07	8.24	.00	0.45	0.73
Caregiver age at intake	0.07	0.10	0.70	.48	-0.13	0.28
Child age at intake	-1.98	2.18	-0.91	.36	-6.26	2.30
Intervention group	-3.76	1.52	-2.47	.01	-6.74	-0.78
Baseline sensitivity WITH						
Baseline depression symptoms	-0.05	0.66	-0.07	.94	-1.34	1.25
Follow-up sensitivity WITH						
Follow-up depression symptoms	-1.29	0.56	-2.31	.02	-2.38	-0.20
Caregiver age at intake WITH						
Baseline sensitivity	1.64	0.50	3.25	.00	0.65	2.62
Baseline depression symptoms	1.47	6.89	0.21	.83	-12.04	14.97

total indirect effect was not significant,  $b = -0.82$ ,  $p = .074$  (see Supplementary Table 3).

### Exploratory Analyses

Finally, given the significant regression models of ABC intervention predicting follow-up levels of sensitivity and parental depression symptoms for ABC completers versus waitlist participants, we conducted a series of exploratory analyses examining potential moderators of these effects. We first attempted to conduct these analyses within the cross-lagged model of completers; however, given the poor fit of the model, all analyses were conducted using PROCESS 2.13 (Hayes, 2014) and pairwise deletion. The results of all of the moderation analyses were consistent between the cross-lagged model and PROCESS.

#### Moderators of intervention effects on sensitivity

First, we examined potential moderators with ABC intervention group as the predictor, follow-up parental sensitivity as the outcome, and baseline sensitivity as a covariate. The addition of an ACE score interaction term to the model did not result in a significant increase in  $R^2$ ,  $\Delta R^2 = .02$ ;  $F(1, 121) = 3.48$ ,  $p = .065$ . Although the moderation was not significant, examining the trend indicated that as baseline ACE score decreased, the effect of intervention on sensitivity increased. Given the nonsignificant moderation effect, these results should be interpreted with caution but suggest that the intervention was most effective in increasing sensitivity for those low in baseline ACE scores (see Table 4). The

addition of a cumulative risk interaction term to the model did not produce a significant change in  $R^2$ , indicating that cumulative risk was not a significant moderator of the association between ABC and sensitivity,  $\Delta R^2 = .00$ ;  $F(1, 125) = 0.49$ ,  $p = .486$ .

#### Moderators of intervention effects on depression

Next, we examined potential moderators with ABC intervention group as the predictor, follow-up parental depression as the outcome, and baseline parental depression as a covariate. The addition of a baseline ACE score interaction term to the model did not produce a significant change in  $R^2$ ,  $\Delta R^2 = .00$ ;  $F(1, 119) = 1.03$ ,  $p = .313$ . Similarly, the addition of a cumulative risk interaction term did not produce a significant change in  $R^2$ ,  $\Delta R^2 = .00$ ,  $F(1, 122) = 0.69$ ,  $p = .409$ . Thus, neither baseline ACE score nor cumulative risk were significant moderators.

### Discussion

The present study replicates and extends previous studies of ABC's effectiveness in contexts of community implementation. Consistent with previous ABC studies reporting improvements in parental sensitivity, ABC participation predicted greater increases in parental sensitivity (i.e., following the child's lead) from baseline to follow-up than assignment to the waitlist control condition. However, ABC did not predict significant baseline to follow-up changes in intrusiveness or positive regard. In addition, parents who completed ABC had greater decreases in baseline to follow-up parental depression symptoms than parents assigned to

**Table 4.** Regression analysis testing the interaction of intervention group and ACE Score on sensitivity

Variable	Unstandardized estimate	SE	t	p	95% CI [lower, upper]
(Constant)	1.62	0.28	5.68	.000	[1.05, 2.18]
Baseline sensitivity	0.46	0.09	4.98	.000	[0.27, 0.64]
ABC	0.67	0.23	2.95	.004	[0.22, 1.12]
Baseline ACE	0.01	0.03	0.29	.769	[-0.06, 0.07]
ABC × Baseline ACE	-0.09	0.05	-1.87	.065	[-0.18, 0.01]
Conditional effect of Intervention Group on Sensitivity for ACE Score					
One SD below mean	0.57	0.19	3.06	.003	[0.20, 0.94]
At mean	0.32	0.13	2.44	.016	[0.06, 0.59]
One SD above mean	0.08	0.19	0.40	.687	[-0.30, 0.45]

Note:  $N = 126$

a waitlist condition. The effect of ABC on parental sensitivity was not mediated by changes in depression (or vice versa). Finally, exploratory analyses indicated that neither cumulative risk nor parental ACEs significantly moderated ABC-related changes in parental sensitivity or depression symptoms.

Although the present study replicates previous findings that ABC is associated with improved parental sensitivity, the lack of changes in intrusiveness and positive regard differ from previous results (Caron et al., 2016; Yarger et al., 2016; Yarger et al., 2019). One prior study examining community implementation of ABC found that in-the-moment feedback from parent coaches was not associated with changes in parental positive regard and suggested that positive regard may be a more difficult intervention target to change than sensitivity and intrusiveness (Caron et al., 2018). Future studies should examine possible explanations for different findings for specific types of parental behavior, such as aspects of ABC implementation (e.g., parent coaches' focus on particular behavior targets) or characteristics of the population receiving ABC.

The influence of ABC on parental depression represents a novel finding. It is possible that weekly visits from parent coaches and receiving positive feedback about their parenting increased parents' feelings of support and competence, thus promoting improvements in depression symptoms, as multiple studies have found evidence that social support is associated with lower depression symptoms in parents (e.g., Bost, Cox, Burchinal, & Payne, 2002; Leahy-Warren, McCarthy, & Corcoran, 2012). In addition, it is possible that changes in parental depression symptoms and sensitivity have bidirectional effects on each other. Although neither parental depression symptoms nor sensitivity were significant mediators, both indirect effects approached statistical significance, suggesting that changes in these parental outcome variables may influence each other. Increased sensitivity may improve parent-child interactions and increase parental self-efficacy, which has been associated with lower parental depression symptoms (Jones & Prinz, 2005). In turn, improvements in depression symptoms may reflect improvements in well-being that allow parents to be more responsive to their children, as previous research indicates that parents' well-being is critical to their ability to support their children effectively (Luthar & Eisenberg, 2017). Future longitudinal studies that measure parental sensitivity and depression at multiple points following ABC will elucidate the directionality of these potential effects.

### Implications for ABC's Effectiveness

These findings provide continued support for ABC's effectiveness in community-based settings. Of particular note, these effects were found in a community organization that implements ABC through the efforts of parent coaches and staff hired primarily from the community being served, highlighting the potential for successful and sustainable dissemination of ABC beyond research-based settings. Furthermore, the lack of significant moderation findings suggests ABC's effectiveness may not be limited to particular subgroups of the population and thus can be applied broadly among community populations. Early intervention has been identified as having a substantial influence on child outcomes as well as the greatest rate of return for resources invested in intervention (Doyle et al., 2009), underlining the importance of continued efforts to disseminate ABC broadly in community settings for improvements both in individual and broader societal outcomes.

The influence of ABC on parental depression symptoms provides additional support for the strength of this intervention. Parental depression is an important early intervention target, given that it is associated with increased distress for the caregiver and a broad range of negative developmental outcomes in the child. For example, having a depressed caregiver in infancy is associated with lower cognitive development (Azak, 2012), higher cortisol levels (Khoury et al., 2016), lower rates of secure attachment (Martins & Gaffan, 2000), and lower social and emotional regulation (Feldman et al., 2009). In early and late childhood, parental depression is further associated with greater psychopathology and emotion regulation difficulties (Goodman et al., 2011), lower social competence and school adjustment (Kersten-Alvarez et al., 2012), reduced emotion regulation (Feng et al., 2008), and emotion and behavior difficulties (van der Waerden et al., 2015). Thus, ABC may promote positive outcomes for children not only through changes in parental sensitivity but also through changes in parental depression.

### Limitations

The design of this study as an RCT was a significant strength relative to previous dissemination studies of ABC that have used pre-post designs. However, these findings were limited by comparison with a waitlist control group rather than with a control

intervention. This design was used to ensure that all families would eventually receive ABC, consistent with Power of Two's goal of disseminating ABC to the community. Waitlist control research designs are common in community-based intervention effectiveness studies (e.g., Caserta *et al.*, 2018; Singla, Kumbakumba, & Aboud, 2015). Nonetheless, future studies should examine ABC efficacy as compared with other evidence-supported parenting interventions, such as Child-Parent Psychotherapy, Parent-Child Interaction Therapy, the Positive Parenting Program, or the Nurse Family Partnership. Identifying shared core components and differences among programs may allow us to match individual families to interventions from which they are most likely to benefit, maximizing effectiveness. Such an approach is consistent with recent moves toward precision home-visiting, in which interventions are tailored to meet families' needs while maintaining fidelity to core active ingredients (Haroz *et al.*, 2019).

A further consideration when interpreting results of the study is attrition both in research participation and intervention completion. Of the 200 dyads initially enrolled in the study, 158 (79%) completed the research follow-up visit. Additionally, of the 100 dyads randomly assigned to receive ABC, 37 (37%) did not complete all 10 sessions of the intervention. This rate of attrition is comparable to those found in other home-based interventions, which often have attrition rates of 20% to 67% (Gomby, 1999). To address this limitation, we conducted both full sample and intervention completer analyses, finding that some effects held across both approaches (i.e., changes in parental sensitivity) and others only for completer analyses (i.e., changes in parental depression).

Finally, an additional limitation was potential variability across staff and participants in attempts to contact participants who cancelled or did not show up for scheduled intervention visits. Consistent with an effectiveness trial, the program's procedures for attempting to reach families receiving the intervention were followed for all participants in the treatment group. Although attempts were made to contact all families after a no show or cancellation, consistent records were not kept, so we were unable to account for this potential variability.

### *Future Directions*

The results of this study suggest many future directions. Examination of these outcomes, particularly the novel depression finding, in diverse populations is critical to ensure consistent effectiveness of ABC as it is implemented in different community settings. The current sample reflected a population with high racial and ethnic minority status, low income, and high levels of risk factors. As a result, future studies will need to examine whether these effectiveness findings generalize to other populations.

Similarly, future studies should continue to examine potential moderators of ABC effectiveness within different populations. Additional suggested moderators include parent characteristics such as substance use, life stressors, psychopathology, and attachment; child characteristics, such as temperament, behavior problems, or prenatal risk; and the interaction between parent and child characteristics (Bernard *et al.*, 2012). Given the associations of parental ACEs and childhood cumulative risk with adverse child outcomes in previous studies (e.g., Evans, Li, & Whipple, 2013; Schickedanz, Halfon, Sastry, & Chung, 2018; Stepleton *et al.*, 2018), moderators should also be examined in relation to ABC's influence on child outcomes. Identifying moderators

would allow organizations to identify parents who are likely to benefit from ABC or who may require additional support.

In addition, future studies should continue to examine mechanisms of treatment effectiveness, including whether parent outcomes such as decreases in depression and improvements in sensitivity mediate ABC's influence on child outcomes. Previous studies have shown that changes in parental sensitivity may be mediators of ABC's effect on child compliance at 36 months of age (Lind *et al.*, 2019) and ABC-T's effect on children's receptive vocabulary at 36 to 60 months of age (Raby *et al.*, 2019). Additional mediation pathways for ABC's influence on parent and child outcomes are likely to be discovered with further research, revealing mechanisms to target during intervention and increasing our empirical knowledge of ABC's effectiveness. In particular, examining such mediation models within the context of RCTs provides a unique opportunity to expand our knowledge of developmental psychology. Random assignment to ABC or waitlist control produces a manipulation of parenting that allows us to test causal pathways between parent and child outcomes in an experimental design. Thus, future studies examining mechanisms of ABC treatment effectiveness have the potential to inform both basic developmental science and applied intervention research.

Given that the present study corroborates the strong empirical basis for ABC's efficacy and effectiveness in promoting positive parent outcomes, an essential future direction is to examine factors that contribute to successful dissemination of ABC, and its developers have identified key components of successful ABC dissemination including selection of sites with the commitment and resources to support ABC fully, screening of potential parent coaches, training on in-the-moment commenting and coding, and supervision that includes both clinical consultation and fidelity monitoring (Dozier & Bernard, 2019). Findings from previous studies support the importance of screening measures for selecting parent coaches who are likely to display strong fidelity to in-the-moment commenting (Caron, Roben, *et al.*, 2018) and of fidelity-focused consultation during implementation for improving coaches' in-the-moment commenting (Caron & Dozier, 2019). Future research can expand on these findings by examining additional variables that contribute to treatment fidelity and successful dissemination. Such variables may include the structure of community organizations; sources of support for dissemination efforts; community engagement; and characteristics of coaches, trainers, and supervisors. Identifying which factors predict successful dissemination is essential to supporting ABC's effectiveness as it is scaled in community settings.

Finally, given the attrition rates, future studies should examine preintervention characteristics that predict likelihood of intervention drop-out (such as baseline parental sensitivity and cumulative risk in the present study) to help identify dyads who may benefit from additional support throughout the intervention. Future studies should also consider what supports may be necessary to facilitate completion of the ABC intervention following enrollment. Previous studies have found that strategies such as engaging community stakeholders, providing additional support to the family, enhancing the program, and training home visitors in motivational interviewing techniques have increased retention in home visiting programs (Biggs, Sprague-Jones, Garstka, & Richardson, 2018; Folger *et al.*, 2016). Power of Two has begun to incorporate a variety of engagement techniques such as holding community events and connecting families to sources of instrumental assistance. An

important future direction will be to test the effectiveness of such efforts empirically.

Overall, this study provides support for ABC's effectiveness and feasibility when implemented as a community-based intervention. These results are promising in supporting ABC as an intervention that may reach underserved families in need, especially given its relatively brief duration and its success in employing interventionists of various backgrounds from within the community. Continued efforts to disseminate ABC on a broad community scale have potential to yield positive outcomes for parents and children alike.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S0954579420000310>.

**Acknowledgments.** We are grateful for the research-practice partnership with Power of Two, specifically with Executive Directors Anne Heller (former) and Erasma Monticciolo (current). We thank the members of the outreach staff and parent coaches at Power of Two who facilitated recruitment and delivered the intervention as well as the participating families.

**Financial Support.** This work was supported by a grant from the Fund for the City of New York to K. Bernard via funding awarded to Power of Two for their ongoing implementation and evaluation of ABC. Direct sources of funding to Power of Two during the evaluation period included: Viola W. Bernard Foundation, Kenworth-Swift Foundation, Marks Family Foundation, Arbor Brothers, The Sirus Fund, Tiger Foundation, Propel Capital, Draper Richards Kaplan, Altman Foundation, Stella & Charles Guttman Foundation, Flom Foundation, Edith Glick Shoolman Children's Foundation, Help for Children/Hedge Funds Care, Brooklyn Branches, ALLINBKLYN, and support from anonymous donors.

## References

- Ainsworth, M. D. S., Bell, S. M., & Stayton, D. J. (1974). Infant-mother attachment and social development: Socialization as a product of reciprocal responsiveness to signals. In M. P. M. Richard (Ed.), *The integration of a child into a social world* (pp. 99-135). London, UK: Cambridge University Press.
- Albers, E. M., Riksen-Walraven, J. M., Sweep, F. C., & de Weerth, C. D. (2008). Maternal behavior predicts infant cortisol recovery from a mild everyday stressor. *Journal of Child Psychology and Psychiatry*, *49*, 97-103. doi:10.1111/j.1469-7610.2007.01818.x
- Azak, S. (2012). Maternal depression and sex differences shape the infants' trajectories of cognitive development. *Infant Behavior and Development*, *35*, 803-814. doi:10.1016/j.infbeh.2012.07.017
- Bagner, D. M., & Graziano, P. A. (2013). Barriers to success in parent training for young children with developmental delay: The role of cumulative risk. *Behavior Modification*, *37*, 356-377. doi:10.1177/0145445512465307
- Bailey, H. N., DeOliveira, C. A., Wolfe, V. V., Evans, E. M., & Hartwick, C. (2012). The impact of childhood maltreatment history on parenting: A comparison of maltreatment types and assessment methods. *Child Abuse & Neglect*, *36*, 236-246. doi:10.1016/j.chiabu.2011.11.005
- Bakermans-Kranenburg, M. J., van IJzendoorn, M. H., & Juffer, F. (2003). Less is more: Meta-analyses of sensitivity and attachment interventions in early childhood. *Psychological Bulletin*, *129*, 195-215. doi:10.1037/0033-2909.129.2.195
- Barnett, M. A., Gustafsson, H., Deng, M., Mills-Koonce, W. R., & Cox, M. (2012). Bidirectional associations among sensitive parenting, language development, and social competence. *Infant and Child Development*, *21*, 374-393. doi:10.1002/icd.1750
- Berlin, L. J., Martoccio, T. L., Bryce, C. I., & Harden, B. J. (2019). Improving infants' stress-induced cortisol regulation through attachment-based intervention: A randomized controlled trial. *Psychoneuroendocrinology*, *103*, 225-232. doi:10.1016/j.psyneuen.2019.01.005
- Berlin, L. J., Martoccio, T. L., & Jones Harden, B. (2018). Improving Early Head Start's impacts on parenting through attachment-based intervention: A randomized controlled trial. *Developmental Psychology*, *54*, 2316-2327. doi:10.1037/dev0000592
- Berlin, L. J., Shanahan, M., & Appleyard Carmody, K. (2014). Promoting supportive parenting in new mothers with substance-use problems: A pilot randomized trial of residential treatment plus an attachment-based parenting program. *Infant Mental Health Journal*, *35*, 81-85. doi:10.1002/imhj.21427
- Bernard, K., Dozier, M., Bick, J., & Gordon, M. K. (2015). Intervening to enhance cortisol regulation among children at risk for neglect: Results of a randomized clinical trial. *Development and Psychopathology*, *27*, 829-841. doi:10.1017/s095457941400073x
- Bernard, K., Dozier, M., Bick, J., Lewis-Morrarty, E., Lindhiem, O., & Carlson, E. (2012). Enhancing attachment organization among maltreated children: Results of a randomized clinical trial. *Child Development*, *83*, 623-636. doi:10.1111/j.1467-8624.2011.01712.x
- Bernard, K., Hostinar, C. E., & Dozier, M. (2015). Intervention effects on diurnal cortisol rhythms of Child Protective Services-referred infants in early childhood: Preschool follow-up results of a randomized clinical trial. *JAMA Pediatrics*, *169*, 112-119. doi:10.1001/jamapediatrics.2014.2369
- Bernard, K., Lee, A. H., & Dozier, M. (2017). Effects of the ABC intervention on foster children's receptive vocabulary: Follow-up results from a randomized clinical trial. *Child Maltreatment*, *22*, 174-179. doi:10.1177/1077559517691126
- Bernard, K., Nissim, G., Vaccaro, S., Harris, J. L., & Lindhiem, O. (2018). Association between maternal depression and maternal sensitivity from birth to 12 months: A meta-analysis. *Attachment & Human Development*, *20*, 578-599. doi:10.1080/14616734.2018.1430839
- Bert, S. C., Guner, B. M., Lanzi, R. G., & Centers for Prevention of Child Neglect (2009). The influence of maternal history of abuse on parenting knowledge and behavior. *Family Relations*, *58*, 176-187. doi:10.1111/j.1741-3729.2008.00545.x
- Bick, J., & Dozier, M. (2013). The effectiveness of an attachment-based intervention in promoting foster mothers' sensitivity toward foster infants. *Infant Mental Health Journal*, *34*, 95-103. doi:10.1002/imhj.21373
- Bick, J., Palmwood, E. N., Zajac, L., Simons, R., & Dozier, M. (2019). Early parenting intervention and adverse family environments affect neural function in middle childhood. *Biological Psychiatry*, *85*, 326-335. doi:10.1016/j.biopsych.2018.09.020
- Biggs, J., Sprague-Jones, J., Garstka, T., & Richardson, D. (2018). Brief motivational interviewing training for home visitors: Results for caregiver retention and referral engagement. *Children and Youth Services Review*, *94*, 56-64. doi:10.1016/j.chilyouth.2018.09.021
- Bost, K. K., Cox, M. J., Burchinal, M. R., & Payne, C. (2002). Structural and supportive changes in couples' family and friendship networks across the transition to parenthood. *Journal of Marriage and Family*, *64*, 517-531. doi:10.1111/j.1741-3737.2002.00517.x
- Caron, E. B., Bernard, K., & Dozier, M. (2018). In vivo feedback predicts parent behavior change in the attachment and biobehavioral catch-up intervention. *Journal of Clinical Child & Adolescent Psychology*, *47*, S35-S46. doi:10.1080/15374416.2016.1141359
- Caron, E. B., & Dozier, M. (2019). Effects of fidelity-focused consultation on clinicians' implementation: An exploratory multiple baseline design. *Administration and Policy in Mental Health and Mental Health Services Research*, *46*, 445-457. doi:10.1007/s10488-019-00924-3
- Caron, E. B., Roben, C. K., Yarger, H. A., & Dozier, M. (2018). Novel methods for screening: Contributions from attachment and biobehavioral catch-up. *Prevention Science*, *19*, 894-903. doi:10.1007/s1121-018-0894-5
- Caron, E. B., Weston-Lee, P., Haggerty, D., & Dozier, M. (2016). Community implementation outcomes of attachment and biobehavioral catch-up. *Child Abuse & Neglect*, *53*, 128-137. doi:10.1016/j.chiabu.2015.11.010
- Caserta, A., Fabiano, G. A., Hulme, K., Pyle, K., Isaacs, L., & Jerome, S. (2018). A waitlist-controlled trial of behavioral parent training for fathers of preschool children. *Evidence-Based Practice in Child and Adolescent Mental Health*, *3*, 106-116. doi:10.1080/23794925.2018.1446768
- Chapman, D. P., Whitfield, C. L., Felitti, V. J., Dube, S. R., Edwards, V. J., & Anda, R. F. (2004). Adverse childhood experiences and the risk of depressive disorders in adulthood. *Journal of Affective Disorders*, *82*, 217-225. doi:10.1016/j.jad.2003.12.013
- Cuijpers, P., Weitz, E., Karyotaki, E., Garber, J., & Andersson, G. (2015). The effects of psychological treatment of maternal depression on children and

- parental functioning: A meta-analysis. *European Child & Adolescent Psychiatry*, 24, 237–245. doi:10.1007/s00787-014-0660-6
- De Wolff, M. S., & van IJzendoorn, M. H. (1997). Sensitivity and attachment: A meta-analysis on parental antecedents of infant attachment. *Child Development*, 68, 571–591. doi:10.2307/1132107
- Doyle, O., Harmon, C. P., Heckman, J. J., & Tremblay, R. E. (2009). Investing in early human development: Timing and economic efficiency. *Economics & Human Biology*, 7, 1–6. doi:10.1016/j.ehb.2009.01.002
- Dozier, M., & Bernard, K. (2017). Attachment and biobehavioral catch-up: Addressing the needs of infants and toddlers exposed to inadequate or problematic caregiving. *Current Opinion in Psychology*, 15, 111–117. doi:10.1016/j.copsyc.2017.03.003
- Dozier, M., & Bernard, K. (2019). *Coaching parents of vulnerable infants: The attachment and biobehavioral catch-up approach*. New York, NY: The Guilford Press.
- Dozier, M., Lindhiem, O., Lewis, E., Bick, J., Bernard, K., & Peloso, E. (2009). Effects of a foster parent training program on young children's attachment behaviors: Preliminary evidence from a randomized clinical trial. *Child and Adolescent Social Work Journal*, 26, 321–332. doi:10.1007/s10560-009-0165-1
- Dozier, M., Peloso, E., Lindhiem, O., Gordon, M. K., Manni, M., Sepulveda, S., . . . Levine, S. (2006). Developing evidence-based interventions for foster children: An example of a randomized clinical trial with infants and toddlers. *Journal of Social Issues*, 62, 767–785. doi:10.1111/j.1540-4560.2006.00486.x
- Dozier, M., Roben, C. K., Caron, E. B., Hoyer, J., & Bernard, K. (2018). Attachment and biobehavioral catch-up: An evidence-based intervention for vulnerable infants and their families. *Psychotherapy Research*, 28, 18–29. doi:10.1080/10503307.2016.1229873
- Dube, S. R., Williamson, D. F., Thompson, T., Felitti, V. J., & Anda, R. F. (2004). Assessing the reliability of retrospective reports of adverse childhood experiences among adult HMO members attending a primary care clinic. *Child Abuse & Neglect*, 28, 729–737. doi:10.1016/j.chiabu.2003.08.009
- Eckenrode, J., Campa, M. I., Morris, P. A., Henderson, C. R., Jr., Bolger, K. E., Kitzman, H., & Olds, D. L. (2017). The prevention of child maltreatment through the Nurse Family Partnership Program: Mediating effects in a long-term follow-up study. *Child Maltreatment*, 22, 92–99. doi:10.1177/1077559516685185
- Evans, G. W., Li, D., & Whipple, S. S. (2013). Cumulative risk and child development. *Psychological Bulletin*, 139, 1342–1396. doi:10.1037/a0031808
- Feldman, R., Granat, A., Pariente, C., Kanety, H., Kuint, J., & Gilboa-Schechtman, E. (2009). Maternal depression and anxiety across the postpartum year and infant social engagement, fear regulation, and stress reactivity. *Journal of the American Academy of Child & Adolescent Psychiatry*, 48, 919–927. doi:10.1097/chi.0b013e3181b21651
- Felitti, V. J., Anda, R. F., Nordenberg, D., Williamson, D. F., Spitz, A. M., Edwards, V., . . . Marks, J. S. (1998). Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventive Medicine*, 14, 245–258. doi:10.1016/s0749-3797(98)00017-8
- Feng, X., Shaw, D. S., Kovacs, M., Lane, T., O'Rourke, F. E., & Alarcon, J. H. (2008). Emotion regulation in preschoolers: The roles of behavioral inhibition, maternal affective behavior, and maternal depression. *Journal of Child Psychology and Psychiatry*, 49, 132–141. doi:10.1111/j.1469-7610.2007.01828.x
- Field, T. (2010). Postpartum depression effects on early interactions, parenting, and safety practices: A review. *Infant Behavior and Development*, 33, 1–6. doi:10.1016/j.infbeh.2009.10.005
- Folger, A. T., Brentley, A. L., Goyal, N. K., Hall, E. S., Sa, T., Peugh, J. L., . . . Ammerman, R. T. (2016). Evaluation of a community-based approach to strengthen retention in early childhood home visiting. *Prevention Science*, 17, 52–61. doi:10.1007/s11121-015-0600-9
- Glasgow, R. E., Lichtenstein, E., & Marcus, A. C. (2003). Why don't we see more translation of health promotion research to practice? Rethinking the efficacy-to-effectiveness transition. *American Journal of Public Health*, 93, 1261–1267. doi:10.2105/ajph.93.8.1261
- Gomby, D. S. (1999). Understanding evaluations of home visitation programs. *Future of Children*, 9, 27–43. doi:10.2307/1602720
- Goodman, S. H., Rouse, M. H., Connell, A. M., Broth, M. R., Hall, C. M., & Heyward, D. (2011). Maternal depression and child psychopathology: A meta-analytic review. *Clinical Child and Family Psychology Review*, 14, 1–27. doi:10.1007/s10567-010-0080-1
- Guild, D. J., Toth, S. L., Handley, E. D., Rogosch, F. A., & Cicchetti, D. (2017). Attachment security mediates the longitudinal association between child–parent psychotherapy and peer relations for toddlers of depressed mothers. *Development and Psychopathology*, 29, 587–600. doi:10.1017/s0954579417000207
- Haltigan, J. D., Roisman, G. I., & Fraley, R. C. (2013). The predictive significance of early caregiving experiences for symptoms of psychopathology through midadolescence: Enduring or transient effects? *Development and Psychopathology*, 25, 209–221. doi:10.1017/s0954579412000260
- Haroz, E. E., Ingalls, A., Kee, C., Goklish, N., Neault, N., Begay, M., & Barlow, A. (2019). Informing precision home visiting: Identifying meaningful subgroups of families who benefit most from Family Spirit. *Prevention Science*, 20, 1244–1254. doi:10.1007/s11121-019-01039-9
- Hayes, A. F. (2014). The PROCESS macro for SPSS (Version 2.13) [Software]. Retrieved from <http://www.afhayes.com>
- Hoagwood, K., & Olin, S. S. (2002). The NIMH blueprint for change report: Research priorities in child and adolescent mental health. *Journal of the American Academy of Child & Adolescent Psychiatry*, 41, 760–767. doi:10.1097/00004583-200207000-00006
- Hu, L. T., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling: A Multidisciplinary Journal*, 6, 1–55. doi:10.1080/10705519909540118
- Jones, T. L., & Prinz, R. J. (2005). Potential roles of parental self-efficacy in parent and child adjustment: A review. *Clinical Psychology Review*, 25, 341–363. doi:10.1016/j.cpr.2004.12.004
- Kersten-Alvarez, L. E., Hosman, C. M. H., Riksen-Walraven, J. M., van Doesum, K. T. M., & Hoefnagels, C. (2011). Which preventive interventions effectively enhance depressed mothers' sensitivity? A meta-analysis. *Infant Mental Health Journal*, 32, 362–376. doi:10.1002/imhj.20301
- Kersten-Alvarez, L. E., Hosman, C. M., Riksen-Walraven, J. M., van Doesum, K. T., Smeekens, S., & Hoefnagels, C. (2012). Early school outcomes for children of postpartum depressed mothers: Comparison with a community sample. *Child Psychiatry & Human Development*, 43, 201–218. doi:10.1007/s10578-011-0257-y
- Khoury, J. E., Gonzalez, A., Levitan, R., Masellis, M., Basile, V., & Atkinson, L. (2016). Maternal self-reported depressive symptoms and maternal cortisol levels interact to predict infant cortisol levels. *Infant Mental Health Journal*, 37, 125–139. doi:10.1002/imhj.21554
- La Greca, A. M., Silverman, W. K., & Lochman, J. E. (2009). Moving beyond efficacy and effectiveness in child and adolescent intervention research. *Journal of Consulting and Clinical Psychology*, 77, 373–382. doi:10.1037/a0015954
- Leahy-Warren, P., McCarthy, G., & Corcoran, P. (2012). First-time mothers: Social support, maternal parental self-efficacy and postnatal depression. *Journal of Clinical Nursing*, 21, 388–397. doi:10.1111/j.1365-2702.2011.03701.x
- Leerkes, E. M., Blankson, A. N., & O'Brien, M. (2009). Differential effects of maternal sensitivity to infant distress and nondistress on social-emotional functioning. *Child Development*, 80, 762–775. doi:10.1111/j.1467-8624.2009.01296.x
- Lengua, L. J., Honorado, E., & Bush, N. R. (2007). Contextual risk and parenting as predictors of effortful control and social competence in preschool children. *Journal of Applied Developmental Psychology*, 28, 40–55. doi:10.1016/j.appdev.2006.10.001
- Lewis-Morrarty, E., Dozier, M., Bernard, K., Terracciano, S. M., & Moore, S. V. (2012). Cognitive flexibility and theory of mind outcomes among foster children: Preschool follow-up results of a randomized clinical trial. *Journal of Adolescent Health*, 51, S17–S22. doi:10.1016/j.jadohealth.2012.05.005
- Lieberman, A. F., Ippen, C. G., & Van Horn, P. (2006). Child-parent psychotherapy: 6-month follow-up of a randomized controlled trial. *Journal of the American Academy of Child & Adolescent Psychiatry*, 45, 913–918. doi:10.1097/01.chi.0000222784.03735.92
- Lieberman, A. F., Van Horn, P., & Ippen, C. G. (2005). Toward evidence-based treatment: Child-parent psychotherapy with preschoolers exposed to marital violence. *Journal of the American Academy of Child & Adolescent Psychiatry*, 44, 1241–1248. doi:10.1097/01.chi.0000181047.59702.58

- Lind, T., Bernard, K., Ross, E., & Dozier, M. (2014). Intervention effects on negative affect of CPS-referred children: Results of a randomized clinical trial. *Child Abuse & Neglect*, 38, 1459–1467. doi:10.1016/j.chiabu.2014.04.004
- Lind, T., Bernard, K., Yarger, H. A., & Dozier, M. (2019). Promoting compliance in children referred to child protective services: A randomized clinical trial. *Child Development*. doi:10.1111/cdev.13207
- Lind, T., Raby, K. L., Caron, E. B., Roben, C. K., & Dozier, M. (2017). Enhancing executive functioning among toddlers in foster care with an attachment-based intervention. *Development and Psychopathology*, 29, 575–586. doi:10.1017/s0954579417000190
- Lovejoy, M. C., Graczyk, P. A., O'Hare, E., & Neuman, G. (2000). Maternal depression and parenting behavior: A meta-analytic review. *Clinical Psychology Review*, 20, 561–592. doi:10.1016/s0272-7358(98)00100-7
- Luthar, S. S., & Eisenberg, N. (2017). Resilient adaptation among at-risk children: Harnessing science toward maximizing salutary environments. *Child Development*, 88, 337–349. doi:10.1111/cdev.12737
- Martins, C., & Gaffan, E. A. (2000). Effects of early maternal depression on patterns of infant–mother attachment: A meta-analytic investigation. *The Journal of Child Psychology and Psychiatry and Allied Disciplines*, 41, 737–746. doi:10.1017/s0021963099005958
- Mesman, J., van IJzendoorn, M. H., & Bakermans-Kranenburg, M. J. (2012). Unequal in opportunity, equal in process: Parental sensitivity promotes positive child development in ethnic minority families. *Child Development Perspectives*, 6, 239–250. doi:10.1111/j.1750-8606.2011.00223.x
- Muthén, L. K., & Muthén, B. O. (2012–2017). *Mplus8* Version 1.5 [Software]. Los Angeles, CA: Muthén & Muthén.
- NICHD (National Institute of Child Health and Human Development) Early Child Care Research Network. (1999) Child care and mother–child interaction in the first three years of life. *Developmental Psychology*, 35, 1399–1413. doi:10.1037/0012-1649.35.6.1399
- NICHD (National Institute of Child Health and Human Development) Early Child Care Research Network. (2003). Early child care and mother–child interaction from 36 months through first grade. *Infant Behavior and Development*, 26, 345–370. doi:10.1016/s0163-6383(03)00035-3
- Pereira, J., Vickers, K., Atkinson, L., Gonzalez, A., Wekerle, C., & Levitan, R. (2012). Parenting stress mediates between maternal maltreatment history and maternal sensitivity in a community sample. *Child Abuse & Neglect*, 36, 433–437. doi:10.1016/j.chiabu.2012.01.006
- Petersen, A. C., Joseph, J., & Feit, M. (Eds.). (2014). *New directions in child abuse and neglect research*. Washington, DC: The National Academies Press.
- Poobalan, A. S., Aucott, L. S., Ross, L., Smith, W. C. S., Helms, P. J., & Williams, J. H. (2007). Effects of treating postnatal depression on mother–infant interaction and child development: Systematic review. *The British Journal of Psychiatry*, 191, 378–386. doi:10.1192/bjp.bp.106.032789
- Prinz, R. J., Sanders, M. R., Shapiro, C. J., Whitaker, D. J., & Lutzker, J. R. (2009). Population-based prevention of child maltreatment: The US Triple P system population trial. *Prevention Science*, 10, 1–12. doi:10.1007/s1121-009-0123-3
- Raby, K. L., Freedman, E., Yarger, H. A., Lind, T., & Dozier, M. (2019). Enhancing the language development of toddlers in foster care by promoting foster parents' sensitivity: Results from a randomized controlled trial. *Developmental Science*, 22. doi:10.1111/desc.12753
- Radloff, L. S. (1977). The CES-D scale: A self-report depression scale for research in the general population. *Applied Psychological Measurement*, 1, 385–401. doi:10.1177/014662167700100306
- Roben, C. K., Dozier, M., Caron, E. B., & Bernard, K. (2017). Moving an evidence-based parenting program into the community. *Child Development*, 88, 1447–1452. doi:10.1111/cdev.12898
- Roberts, R. E. (1980). Reliability of the CES-D scale in different ethnic contexts. *Psychiatry Research*, 2, 125–134. doi:10.1016/0165-1781(80)90069-4
- Schickedanz, A., Halfon, N., Sastry, N., & Chung, P. J. (2018). Parents' adverse childhood experiences and their children's behavioral health problems. *Pediatrics*, 142. doi:10.1542/peds.2018-0023
- Sethna, V., Murray, L., Netsi, E., Psychogiou, L., & Ramchandani, P. G. (2015). Paternal depression in the postnatal period and early father–infant interactions. *Parenting*, 15, 1–8. doi:10.1080/15295192.2015.992732
- Singla, D. R., Kumbakumba, E., & Aboud, F. E. (2015). Effects of a parenting intervention to address maternal psychological wellbeing and child development and growth in rural Uganda: A community-based, cluster-randomised trial. *The Lancet Global Health*, 3, e458–e469. doi:10.1016/s2214-109x(15)00099-6
- Spoth, R., Redmond, C., Shin, C., Lepper, H., Haggerty, K., & Wall, M. (1998). Risk moderation of parent and child outcomes in a preventive intervention: A test and replication. *American Journal of Orthopsychiatry*, 68, 565–579. doi:10.1037/h0080365
- Stempleton, K., Bosk, E. A., Duron, J. F., Greenfield, B., Ocasio, K., & MacKenzie, M. J. (2018). Exploring associations between maternal adverse childhood experiences and child behavior. *Children and Youth Services Review*, 95, 80–87. doi:10.1016/j.childyouth.2018.10.027
- Stolk, M. N., Mesman, J., van Zeijl, J., Alink, L. R., Bakermans-Kranenburg, M. J., van IJzendoorn, M. H., . . . Koot, H. M. (2008). Early parenting intervention: Family risk and first-time parenting related to intervention effectiveness. *Journal of Child and Family Studies*, 17, 55–83. doi:10.1007/s10826-007-9136-3
- Tabachnick, A. R., Raby, K. L., Goldstein, A., Zajac, L., & Dozier, M. (2019). Effects of an attachment-based intervention in infancy on children's autonomic regulation during middle childhood. *Biological Psychology*, 143, 22–31. doi:10.1016/j.biopsycho.2019.01.006
- Thomas, R., & Zimmer-Gembeck, M. J. (2011). Accumulating evidence for parent–child interaction therapy in the prevention of child maltreatment. *Child Development*, 82, 177–192. doi:10.1111/j.1467-8624.2010.01548.x
- Tømmerås, T., & Kjøbli, J. (2017). Family resources and effects on child behavior problem interventions: A cumulative risk approach. *Journal of Child and Family Studies*, 26, 2936–2947. doi:10.1007/s10826-017-0777-6
- Trentacosta, C. J., Hyde, L. W., Shaw, D. S., Dishion, T. J., Gardner, F., & Wilson, M. (2008). The relations among cumulative risk, parenting, and behavior problems during early childhood. *Journal of Child Psychology and Psychiatry*, 49, 1211–1219. doi:10.1111/j.1469-7610.2008.01941.x
- Valcan, D. S., Davis, H., & Pino-Pasternak, D. (2018). Parental behaviours predicting early childhood executive functions: A meta-analysis. *Educational Psychology Review*, 30, 607–649. doi:10.1007/s10648-017-9411-9
- van der Waerden, J., Galéra, C., Larroque, B., Saurel-Cubizolles, M. J., Sutter-Dallay, A. L., Melchior, M., & EDEN Mother–Child Cohort Study Group. (2015). Maternal depression trajectories and children's behavior at age 5 years. *The Journal of Pediatrics*, 166, 1440–1448. doi:10.1016/j.jpeds.2015.03.002
- Weisz, J. R., Weiss, B., & Donenberg, G. R. (1992). The lab versus the clinic: Effects of child and adolescent psychotherapy. *American Psychologist*, 47, 1578–1585. doi:10.1037/0003-066x.47.12.1578
- Yarger, H. A., Bernard, K., Caron, E. B., Wallin, A., & Dozier, M. (2019). Enhancing parenting quality for young children adopted internationally: Results of a randomized controlled trial. *Journal of Clinical Child & Adolescent Psychology*. doi:10.1080/15374416.2018.1547972
- Yarger, H. A., Hoye, J. R., & Dozier, M. (2016). Trajectories of change in attachment and biobehavioral catch-up among high-risk mothers: A randomized clinical trial. *Infant Mental Health Journal*, 37, 525–536. doi:10.1002/imhj.21585
- Zeegers, M. A., Colonna, C., Stams, G. J. J., & Meins, E. (2017). Mind matters: A meta-analysis on parental mentalization and sensitivity as predictors of infant–parent attachment. *Psychological Bulletin*, 143, 1245–1272. doi:10.1037/bul0000114